


# CS 188: Artificial Intelligence

## Bayes' Nets: Independence



Instructors: Pieter Abbeel & Dan Klein --- University of California, Berkeley

# Probability Recap

- Conditional probability

$$P(x|y) = \frac{P(x, y)}{P(y)}$$

- Product rule

$$P(x, y) = P(x|y)P(y)$$

- Chain rule

$$\begin{aligned} P(X_1, X_2, \dots, X_n) &= P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots \\ &= \prod_{i=1}^n P(X_i|X_1, \dots, X_{i-1}) \end{aligned}$$

- X, Y independent if and only if:  $\forall x, y : P(x, y) = P(x)P(y)$


- X and Y are conditionally independent given Z if and only if:

$$\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$$

$$X \perp\!\!\!\perp Y | Z$$

# Bayes' Nets

- A Bayes' net is an efficient encoding of a probabilistic model of a domain



○ = unobserved  
⊖ = obs. conditioned on

- Questions we can ask:

- Inference: given a fixed BN, what is  $P(X | e)$ ?
- Representation: given a BN graph, what kinds of distributions can it encode?
- Modeling: what BN is most appropriate for a given domain?




# Bayes' Net Semantics

- A directed, acyclic graph, one node per random variable
- A conditional probability table (CPT) for each node
  - A collection of distributions over  $X$ , one for each combination of parents' values

$$P(X|a_1 \dots a_n)$$

- Bayes' nets implicitly encode joint distributions
  - As a product of local conditional distributions
  - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:


$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$





# Example: Alarm Network


B	P(B)
+b	0.001
-b	0.999



E	P(E)
+e	0.002
-e	0.998

A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95

A	M	P(M A)
+a	+m	0.7
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99




B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999


$$P(+b, -e, +a, -j, +m) =$$

# Example: Alarm Network

B	<u>P(B)</u>
+b	<u>0.001</u>
-b	0.999



E	<u>P(E)</u>
+e	<u>0.002</u>
-e	0.998



A	J	<u>P(J A)</u>
+a	+j	0.9
+a	-j	<u>0.1</u>
-a	+j	0.05
-a	-j	0.95

A	M	<u>P(M A)</u>
+a	+m	<u>0.7</u>
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99

B	E	A	<u>P(A B,E)</u>
+b	+e	+a	0.95
+b	+e	-a	0.05
<u>+b</u>	<u>-e</u>	<u>+a</u>	<u>0.94</u>
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999

$$\begin{aligned}
 &P(\underline{+b}, \underline{-e}, \underline{+a}, \underline{-j}, \underline{+m}) = \\
 &P(\underline{+b})P(\underline{-e})P(\underline{+a} | \underline{+b}, \underline{-e})P(\underline{-j} | \underline{+a})P(\underline{+m} | \underline{+a}) = \\
 &\underline{0.001 \times 0.998 \times 0.94 \times 0.1 \times 0.7}
 \end{aligned}$$

$$P(X_1, X_2, X_3)$$

$(M-1) \times M \times M$

each take M states

# Size of a Bayes' Net

- How big is a joint distribution over N Boolean variables?

$$2^N - 1$$



- How big is an N-node net if nodes have up to k parents?

$$O(N * 2^{k+1})$$

- Both give you the power to calculate

$$P(X_1, X_2, \dots, X_n)$$

- BNs: Huge space savings!
- Also easier to elicit local CPTs
- Also faster to answer queries (coming)



# Bayes' Nets

---

- ✓ Representation
  - Conditional Independences
  - Probabilistic Inference
  - Learning Bayes' Nets from Data

# Conditional Independence

- X and Y are **independent** if


$$\forall x, y \quad P(x, y) = P(x)P(y) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y$$

- X and Y are **conditionally independent** given Z

$$\forall x, y, z \quad \underline{P(x, y|z) = P(x|z)P(y|z)} \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y | Z$$

- (Conditional) independence is a property of a distribution

- Example:  $Alarm \perp\!\!\!\perp Fire | Smoke$




# Bayes Nets: Assumptions


- Assumptions we are required to make to define the Bayes net when given the graph:

$$P(x_i | x_1 \cdots x_{i-1}) = P(x_i | \text{parents}(X_i))$$

- Beyond above “chain rule → Bayes net” conditional independence assumptions
  - Often additional conditional independences
  - They can be read off the graph
- Important for modeling: understand assumptions made when choosing a Bayes net graph



# Example




- Conditional independence assumptions directly from simplifications in chain rule:

$$p(x_i | x_{-i}) = \phi(x_i | \text{parents}(x_i))$$

- Additional implied conditional independence assumptions?

# Independence in a BN


- Important question about a BN:
  - Are two nodes independent given certain evidence?
  - If yes, can prove using algebra (tedious in general)
  - If no, can prove with a counter example
  - Example:



- Question: are X and Z necessarily independent?
  - Answer: no. Example: low pressure causes rain, which causes traffic.
  - X can influence Z, Z can influence X (via Y)
  - Addendum: they *could* be independent: how?



# D-separation: Outline



# D-separation: Outline

---

- Study independence properties for triples
- Analyze complex cases in terms of member triples
- D-separation: a condition / algorithm for answering such queries

# Causal Chains

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Guaranteed X independent of Z ? *No!*

- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

- Example:

- Low pressure causes rain causes traffic, high pressure causes no rain causes no traffic

- In numbers:


$$P(+y \mid +x) = 1, P(-y \mid -x) = 1, \\ P(+z \mid +y) = 1, P(-z \mid -y) = 1$$



# Causal Chains

- This configuration is a “causal chain”


- Guaranteed X independent of Z given Y?



$$P(x, y, z) = P(x)P(y|x)P(z|y)$$



$$\begin{aligned}
 P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} \\
 &= \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)} \\
 &= P(z|y)
 \end{aligned}$$




Yes!

- Evidence along the chain “blocks” the influence

# Common Cause

- This configuration is a “common cause”

- Guaranteed X independent of Z? **No!**



- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

- Example:

- Project due causes both forums busy and lab full

- In numbers:


$$\begin{aligned} P(+x \mid +y) &= 1, P(-x \mid -y) = 1, \\ P(+z \mid +y) &= 1, P(-z \mid -y) = 1 \end{aligned}$$

$$\underline{P(x, y, z) = P(y)P(x|y)P(z|y)}$$

# Common Cause

- This configuration is a “common cause”

- Guaranteed X and Z independent given Y?




$$P(x, y, z) = P(y)P(x|y)P(z|y)$$

$$\begin{aligned} \underline{P(z|x, y)} &= \frac{P(x, y, z)}{P(x, y)} \\ &= \frac{\cancel{P(y)}\cancel{P(x|y)}P(z|y)}{\cancel{P(y)}\cancel{P(x|y)}} \\ &= \underline{P(z|y)} \\ &\text{Yes!} \end{aligned}$$

- Observing the cause blocks influence between effects.


# Common Effect

- Last configuration: two causes of one effect (v-structures)



Collider  
explaining

- Are X and Y independent? | nothing
  - **Yes**: the ballgame and the rain cause traffic, but they are not correlated
  - Still need to prove they must be (try it!)
- Are X and Y independent given Z?
  - **No**: seeing traffic puts the rain and the ballgame in competition as explanation.
- This is backwards from the other cases
  - Observing an effect **activates** influence between possible causes.



# The General Case


---






# The General Case

- General question: in a given BN, are two variables independent (given evidence)?
- Solution: analyze the graph
- Any complex example can be broken into repetitions of the three canonical cases



# Reachability

- Recipe: shade evidence nodes, look for paths in the resulting graph
- Attempt 1: if two nodes are connected by an undirected path not blocked by a shaded node, they are conditionally independent
- Almost works, but not quite
  - Where does it break?
  - Answer: the v-structure at T doesn't count as a link in a path unless "active"





# Active / Inactive Paths

Question: Are  $X$  and  $Y$  conditionally independent given evidence variables  $\{Z\}$ ?

- Yes, if  $X$  and  $Y$  “d-separated” by  $Z$
- Consider all (undirected) paths from  $X$  to  $Y$
- No active paths = independence!

directed







A path is active if each triple is active:




- Causal chain  $A \rightarrow B \rightarrow C$  where  $B$  is unobserved (either direction)
- Common cause  $A \leftarrow B \rightarrow C$  where  $B$  is unobserved
- Common effect (aka v-structure)  
 $A \rightarrow B \leftarrow C$  where  $B$  or one of its descendants is observed

All it takes to block a path is a single inactive segment

not blocked  
Active Triples



blocked  
Inactive Triples




# D-Separation

- Query:  $X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\} ?$
- Check all (undirected!) paths between  $X_i$  and  $X_j$ 
  - If one or more active, then independence not guaranteed

$$X_i \not\perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$

- Otherwise (i.e. if all paths are inactive),  
then independence is guaranteed

$$X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$




# Example

---

$R \perp\!\!\!\perp B$       *Yes*

$R \perp\!\!\!\perp B | T$

$R \perp\!\!\!\perp B | T'$



# Example



$P(B|A)$

	B	1
A	0	.5
1	.5	.5

independent  
B

	D	1
A	0	.49
1	.52	.48

depend

$L \perp\!\!\!\perp T' | T$  Yes


$L \perp\!\!\!\perp B$  Yes

$L \perp\!\!\!\perp B | T$  "No"

$L \perp\!\!\!\perp B | T'$  No

$L \perp\!\!\!\perp B | T, R$  Yes

$L \perp\!\!\!\perp B | R$




# Example

- Variables:
  - R: Raining
  - T: Traffic
  - D: Roof drips
  - S: I'm sad
- Questions:

$$\underline{T \perp\!\!\!\perp D}$$

$$T \perp\!\!\!\perp D | R \quad \text{Yes}$$

$$T \perp\!\!\!\perp D | R, S \quad \text{No}$$




$\perp$   
=  
 $\perp\!\!\!\perp$

# Structure Implications

- Given a Bayes net structure, can run d-separation algorithm to build a complete list of conditional independences that are necessarily true of the form

$$X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$



- This list determines the set of probability distributions that can be represented





# Computing All Independences



COMPUTE ALL THE  
INDEPENDENCES!




# Topology Limits Distributions

- Given some graph topology  $G$ , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)
- Adding arcs increases the set of distributions, but has several costs
- Full conditioning can encode any distribution


$$\{X \perp\!\!\!\perp Y, X \perp\!\!\!\perp Z, Y \perp\!\!\!\perp Z, \\ X \perp\!\!\!\perp Z \mid Y, X \perp\!\!\!\perp Y \mid Z, Y \perp\!\!\!\perp Z \mid X\}$$



$$\{X \perp\!\!\!\perp Z \mid Y\}$$



$$\{\}$$



# Bayes Nets Representation Summary

---

- Bayes nets compactly encode joint distributions

*graphical model*

- Guaranteed independencies of distributions can be deduced from BN graph structure

- D-separation gives precise conditional independence guarantees from graph alone

- A Bayes' net's joint distribution may have further (conditional) independence that is not detectable until you inspect its specific distribution

# Bayes' Nets

---

✓ Representation

✓ Conditional Independences

- Probabilistic Inference

- Enumeration (exact, exponential complexity)

- Variable elimination (exact, worst-case exponential complexity, often better)

- Probabilistic inference is NP-complete

- Sampling (approximate)

- Learning Bayes' Nets from Data